



EUROPEAN MEDICINES AGENCY
SCIENCE MEDICINES HEALTH

20 September 2016
EMA/HMPC/166517/2015
Committee on Herbal Medicinal Products (HMPC)

European Union herbal monograph on *Origanum majorana* L., herba

Final

Discussion in Working Party on European Union monographs and list (MLWP)	November 2014 January 2015 March 2015 May 2015 July 2015 November 2015
Adoption by Committee on Herbal Medicinal Products (HMPC) for release for consultation	02 February 2016
Start of public consultation	15 February 2016
End of consultation (deadline for comments ¹)	15 May 2016
Re-discussion in MLWP	May/June 2016
Adoption by HMPC	20 September 2016

Keywords	Herbal medicinal products; HMPC; European Union herbal monographs; traditional use; <i>Origanum majorana</i> L., herba; <i>Origani majoranae</i> herba; Marjoram
----------	--

¹ No comments were received during the period of public consultation. Therefore the final monograph is published together with the final assessment report and list of references, without an 'Overview of comments received during the public consultation'.



BG (bulgarski): Майорана, стрък	LT (lietuvių kalba): Kvapiųjų mairūnų žolė
CS (čeština): dobromyslová nať	LV (latviešu valoda): Majorāna laksts
DA (dansk): Havemerian	MT (Malti): ħaxixa tal-Merdqux
DE (Deutsch): Majorankraut	NL (Nederlands): Echte marjolein, kruid
EL (elliniká): Πόα ορίγανον το αμάρακον	PL (polski): Ziele majeranku
EN (English): Marjoram	PT (português): Manjerona
ES (español): Mejorana, partes aéreas de	RO (română): iarbă de măghiran
ET (eesti keel): majoraaniürt	SK (slovenčina): Vňať majoránu záhradného
FI (suomi): maustemeirami, verso	SL (slovenščina): zel majarona
FR (français): Marjolaine (sommité fleurie de)	SV (svenska): Mejram, ört
HR (hrvatski): mažuranova zelen	IS (íslenska):
HU (magyar): majoránna virágos hajtás	NO (norsk): Merian
IT (italiano): Maggiorana parti aeree fiorite	

European Union herbal monograph on *Origanum majorana* L., herba

1. Name of the medicinal product

To be specified for the individual finished product.

2. Qualitative and quantitative composition^{2,3}

Well-established use	Traditional use
	<p>With regard to the registration application of Article 16d(1) of Directive 2001/83/EC</p> <p><i>Origanum majorana</i>, herba (Majoram)</p> <p>i) Herbal substance</p> <p>Not applicable</p> <p>ii) Herbal preparations</p> <p>Comminuted herbal substance</p> <p>Extract⁴ (ratio of herbal substance to extraction solvent 1:5), extraction solvents ethanol 96% V/V and white petroleum jelly.</p>

3. Pharmaceutical form

Well-established use	Traditional use
	<p>Comminuted herbal substance as herbal tea for oral use.</p> <p>Herbal preparations in semi-solid dosage forms for cutaneous use.</p> <p>The pharmaceutical form should be described by the European Pharmacopoeia full standard term.</p>

² The declaration of the active substance(s) for an individual finished product should be in accordance with relevant herbal quality guidance.

³ The material complies with the monograph Majoranae herba in Farmakopea Polska X, 2014.

⁴ The preparation is described in the Farmakopea Polska (1995), two parts of comminuted *Origanum majorana* L., herba is moistened with one part of ethanol 96% and then warm extracted with ten parts of white petroleum jelly until ethanol evaporation.

4. Clinical particulars

4.1. Therapeutic indications

Well-established use	Traditional use
	<p>Indication 1)</p> <p>Traditional herbal medicinal product used for the symptomatic relief of mild spasmodic gastro-intestinal complaints such as bloating and flatulence.</p> <p>Indication 2)</p> <p>Traditional herbal medicinal product used for relief of irritated skin around the nostrils.</p> <p>The product is a traditional herbal medicinal product for use in the specified indications exclusively based upon long-standing use.</p>

4.2. Posology and method of administration⁵

Well-established use	Traditional use
	<p>Posology</p> <p>Indication 1)</p> <p><i>Adults and elderly</i></p> <p>a) Herbal tea: 2-4 g of the comminuted herbal substance in 150 ml of boiling water as a herbal infusion, once to twice daily before meal.</p> <p>Daily dose: 2-8 g</p> <p>The use in children and adolescents under 18 years of age is not recommended (see section 4.4 'Special warnings and precautions for use').</p> <p>Indication 2)</p> <p>b) <i>Children 1-11 years, adolescents and adults</i></p> <p>Small amount of the preparation spread around nostrils, 2 to 4 times daily</p> <p>The use in children under 1 year of age is not recommended (see section 4.4 'Special warnings and precautions for use').</p>

⁵ For guidance on herbal substance/herbal preparation administered as herbal tea or as infusion/decoction/macerate preparation, please refer to the HMPC 'Glossary on herbal teas' (EMA/HMPC/5829/2010 Rev.1).

Well-established use	Traditional use
	<p>Duration of use</p> <p>Indication 1)</p> <p>If the symptoms persist longer than 2 weeks during the use of the medicinal product, a doctor or a qualified health care practitioner should be consulted.</p> <p>Indication 2)</p> <p>If the symptoms persist longer than 1 week during the use of the medicinal product, a doctor or a qualified health care practitioner should be consulted</p> <p>Method of administration</p> <p><i>Preparation a)</i></p> <p>Oral use</p> <p><i>Preparation b)</i></p> <p>Cutaneous use</p>

4.3. Contraindications

Well-established use	Traditional use
	Hypersensitivity to the active substance or to other plants of the Lamiaceae family.

4.4. Special warnings and precautions for use

Well-established use	Traditional use
	<p>Indication 1)</p> <p>The use in children and adolescents under 18 years of age has not been established due to lack of adequate data.</p> <p>Indication 2)</p> <p>The use in children under 1 year of age has not been established due to lack of adequate data.</p> <p>The deep penetration of the ointment inside nostril should be avoided, as it can reduce the activity of the ciliary epithelium.</p> <p>If signs of skin infection are observed during the use of the medicinal product, a doctor or a qualified health care practitioner should be</p>

Well-established use	Traditional use
	<p>consulted.</p> <p>Eye contact with unwashed hands after the application may potentially cause irritation.</p> <p>Indication 1) and 2)</p> <p>If the symptoms worsen during the use of the medicinal product, a doctor or a qualified health care practitioner should be consulted.</p>

4.5. Interactions with other medicinal products and other forms of interaction

Well-established use	Traditional use
	None reported

4.6. Fertility, pregnancy and lactation

Well-established use	Traditional use
	<p>Safety during pregnancy and lactation has not been established. In the absence of sufficient data, the use during pregnancy and lactation is not recommended.</p> <p>No fertility data available.</p>

4.7. Effects on ability to drive and use machines

Well-established use	Traditional use
	No studies on the ability to drive or use machines have been performed.

4.8. Undesirable effects

Well-established use	Traditional use
	<p>None known</p> <p>If adverse reactions occur, a doctor or a qualified health care practitioner should be consulted.</p>

4.9. Overdose

Well-established use	Traditional use
	No case of overdose has been reported.

5. Pharmacological properties

5.1. Pharmacodynamic properties

Well-established use	Traditional use
	Not required as per Article 16c(1)(a)(iii) of Directive 2001/83/EC.

5.2. Pharmacokinetic properties

Well-established use	Traditional use
	Not required as per Article 16c(1)(a)(iii) of Directive 2001/83/EC.

5.3. Preclinical safety data

Well-established use	Traditional use
	Not required as per Article 16c(1)(a)(iii) of Directive 2001/83/EC, unless necessary for the safe use of the product. Tests on reproductive toxicity, genotoxicity and carcinogenicity have not been performed.

6. Pharmaceutical particulars

Well-established use	Traditional use
	Not applicable

7. Date of compilation/last revision

20 September 2016